

AMINOGRO 13-0-0



GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)..... 13%

Amino Acids..... 80%

DERIVED FROM: Fermented Amino Acid Powder

Aminogro 13-0-0 is a completely solubilized, hydrolyzed, vegetable protein which supports plant growth, reduces abiotic stress and benefits soil health. Aminogro 13-0-0 is completely water soluble allowing the efficient uptake of nitrogen for all your crops. It can be applied as a foliar spray, to the soil, or in your hydroponic tank. Aminogro 13-0-0 contains over 20 beneficial amino acids and peptides and such fermentation by-products as enzymes, vitamins and biostimulants. Frequent applications at lower rates are generally more effective than fewer applications at higher rates.

FOLIAR APPLICATION RATES:

Apply with sufficient water for coverage

CROP	APPLICATION RATES	APPLICATION TIMING
In all crops:	1.5-2 kg/100L/ha	In case of stress, several applications with smaller application rates increase efficiency. 300-500 ml per 100 L water with plant protection treatments
Vegetables such as: Fruiting vegetables, cucurbits, brassica or cole crops; leafy vegetables	2-3 kg/ha	Apply first application 10-14 days after transplanting, then apply 3-4 times at 10-14 day intervals from early flowering and during fruit sizing
Field crops such as: Corn, potatoes, barley, canola, soybeans, sorghum, sugar beets, wheat, lentils, beans, peas	2-3 kg/100L/ha	Apply mid-vegetative and early flowering. Can be mixed with herbicides (including glyphosate) and fungicides. Applications may be made every 10-14 days
Forage crops such as: Alfalfa, ryegrass, timothy, fescue	2-3 kg/100L/ha	Apply when plants are 3-6 inches high before first cut, and then 10-14 days before each cutting.
Tree, fruit and vine crops such as: Apples, cherries, pear, stone fruit, table and wine grapes, and berry crops such as: Blueberries, cranberries, raspberries, strawberries.	2-3 kg/ha	Apply every 10-14 days. Make application from early flowering to early ripening.
Ornamental plants (open field)	2-3 kg/ha	Apply using a 10 day interval.
Greenhouse and Potted Plants – all crops and ornamentals	1-2 kg/ha, 20-40 g/100m ²	Apply using a 10-14 day interval.
Turf	2-4 kg/ha, 20-40 g/100m ²	Make applications beginning after green up. Can be mixed with herbicides, fertilizers, and fungicides. Applications may be made every 10-14 days.

FERTIGATION (DRIP/TRICKLE/SPRINKLER IRRIGATION) AND IN-FURROW/SIDE DRESS APPLICATION RATES:

Dissolve 1 Part Aminogro 13-0-0 to 2 Parts Water and Apply as a Concentrate In-Furrow or Inject into Fertigation System

CROP	APPLICATION RATES	APPLICATION TIMING
Berries such as: Blueberries, cranberries, raspberries, strawberries	2-4 kg/ha	Begin applications approximately 14 days after transplanting and continue until early fruit ripening.
Vegetable such as: Fruiting vegetables, cucurbits, brassica or cole crops	2-3 kg/ha	Begin applications 14-21 days after transplanting. Continue applications on a 14-21 day interval during fruit set and fruit sizing.
Hydroponics and Cannabis	2 g/L	Throughout growth cycle.



ATTENTION

Keep out of reach of children and animals.

Product contains soy. Adverse reactions may occur in sensitive persons. Keep out of reach of children and animals. Wear dust mask and protective gloves. If skin contact occurs, wash with soap and water. If allergic reaction occurs, seek medical attention. Store in a dry, well ventilated location. Dispose of contents and container in accordance with local, regional, and national regulations.

Lot Number:

Expiry Date:

20 KG